

Detection and attribution of human influence on regional precipitation

Article

Accepted Version

Balan Sarojini, B., Stott, P. A. and Black, E. (2016) Detection and attribution of human influence on regional precipitation. *Nature Climate Change*, 6 (7). pp. 669-675. ISSN 1758-678X doi: <https://doi.org/10.1038/nclimate2976> Available at <https://centaur.reading.ac.uk/65605/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1038/nclimate2976>

Publisher: Nature Publishing Group

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Detection and Attribution of Human Influence on Regional Precipitation

Beena Balan Sarojini^{1, 2, 3*}, Peter A. Stott⁴ and Emily Black^{1, 2}

¹National Centre for Atmospheric Science - Climate Directorate, Reading, UK

²Department of Meteorology/Walker Institute, University of Reading, UK

³Department of Geography and Environmental Science, University of Reading, UK

⁴Met Office Hadley Centre, Exeter, UK

25 February 2016
(Revised)

*Corresponding author: Beena Balan Sarojini (b.balansarojini@reading.ac.uk)

NCAS - Climate, Department of Meteorology

University of Reading, Reading RG6 6BB, United Kingdom

Phone: + 44 118 378 6238, Fax: + 44 118 378 8316

Understanding how human influence on climate is affecting precipitation around the world is immensely important for defining mitigation policies, and for adaptation planning. Yet despite increasing evidence for the influence of climate change on global patterns of precipitation, and expectations that significant changes in regional precipitation should have already occurred as a result of human influence on climate, compelling evidence of anthropogenic fingerprints on regional precipitation is obscured by observational and modelling uncertainties and is likely to remain so using current methods for years to come. This is in spite of substantial ongoing improvements in models, new reanalyses and a satellite record that spans over thirty years. If we are to quantify how human-induced climate change is affecting the regional water cycle, we need to consider novel ways of identifying the effects of natural and anthropogenic influences on precipitation that take full advantage of our physical expectations.

How rainfall is changing in a particular region is a question of great practical importance to societies. Floods and droughts threaten the lives and livelihoods of many people and enhancing their resilience is of major concern, particularly as anthropogenic climate change is expected to increase the frequency of floods and droughts¹. These expected changes may, moreover, render risk assessments based purely on the historical record inaccurate. Well-planned adaptation to climate change thus requires information on how hazardous rainfall is changing in response to anthropogenic forcing. Are we observing systematic changes or are we simply experiencing natural variability? This is the business of detection and attribution (Box 1).

62 New observations and improved models have enabled the detection of anthropogenic
63 change in the water cycle at large spatial scales^{2, 3,4}, although even here large
64 uncertainties remain. The Intergovernmental Panel on Climate Change⁵ (IPCC) in its
65 Fifth Assessment Report (AR5) concludes that it is *likely* that anthropogenic
66 influences have affected the global water cycle since 1960. In Section TS. 6.3 of AR5,
67 two key uncertainties which limit confidence in attribution assessments of the causes
68 of precipitation changes are recognised as 1) observational and modelling
69 uncertainties, and 2) the large effect of internal variability. Hence there is only
70 *medium confidence* that there is an anthropogenic contribution to global-scale changes
71 in precipitation patterns over land since 1950, with higher levels of confidence
72 precluded by uncertainty in models and observations and the large internal variability
73 in precipitation⁶.

74 At continental scales, there has been some limited success in detecting anthropogenic
75 changes in land precipitation. Anthropogenically driven changes in zonal averages of
76 land precipitation were detected by e.g., ref. 7 – although in some cases the results
77 were found to be sensitive to the observational dataset used. Anthropogenic trends in
78 precipitation have also been detected in the northern mid-to-high latitude lands^{8,9} and
79 southwest Australia¹⁰, where in both regions there are large expected trends that are
80 coherent over wide areas (Figure TS.16 of IPCC, 2013). In general, however,
81 detection and attribution of an anthropogenic signal at these scales is hampered by
82 observational uncertainty and model error^{2, 6,8,9,11}. Even the continental-scale studies
83 described above are too coarse to inform assessments of the extent to which human-
84 induced climate change has affected changes affecting many people locally. Because
85 internal variability in precipitation tends to increase with reducing spatial scale there
86 may be a tendency to assume that detection of an anthropogenic signal of change is

more likely at global or continental scales than at regional scales. In this context, by regional scales we refer to smaller spatial scales than ‘continental’, typically thinking of areas of the globe characterised by specific geographic and climatological features⁵.

This perspective argues that analysis of changes in the processes governing internal variability in precipitation should facilitate the detection and attribution of anthropogenic changes at a range of spatial scales. In some cases an anthropogenic signal may be easier to detect at regional scales, where we have a clearer expectation of forced changes^{8,9,10}. Above all progress in detection and attribution of changes in the water cycle requires the development of novel metrics, which should help facilitate the identification of significant changes in precipitation even in the presence of substantial modelling and observational uncertainty¹². This should enable faster progress to be made than would be possible by simply waiting for models or observations to improve or by simply waiting for the signal of climate change to strengthen sufficiently to emerge from the noise of internal variability.

We first compare physical expectations of global and regional anthropogenic changes in precipitation. Next, we describe how spatial scale modifies the impact of model error and observational uncertainty on detection of these changes. We then consider how novel methods of analysis can be brought to bear on detection and attribution of regional changes in precipitation. Finally, we reflect on how our current models and observations can best be utilised to provide a robust view of anthropogenic change in regional precipitation.

Expected changes on global and continental scales

Based on the physical relation of Clausius-Clapeyron, surface warming is expected to result in an increase in water vapour concentrations at a rate of 6-7% per Kelvin¹³, given that the relative humidity is expected to remain nearly constant¹⁴. This thermodynamic expectation of an intensification of the water cycle has been confirmed in changes in observed and simulated atmospheric moisture content over land and ocean^{15, 16,17,18}, albeit in observations from recent years there is some evidence of a reduction of relative humidity over land¹⁹.

Global mean precipitation is not, however, expected to scale with the increase in atmospheric moisture because it is controlled not by specific humidity, but by the energy budget of the troposphere. The two complementary energy budget arguments are 1) the tropospheric latent heating during precipitation formation is balanced by the radiative cooling to outer space¹⁴, and 2) at the surface the latent heat flux (which is proportional to global mean evaporation and hence global mean precipitation) is balanced by the sensible and radiative heat fluxes^{14,13,15,20}. The warming of the troposphere increases the radiative cooling rate and hence the precipitation. However, if the warming is driven by an increase in greenhouse gases (GHGs), the increase in the radiative cooling rate is partly offset by the direct radiative effect of the GHGs, which is to decrease the radiative cooling rate. This implies that the precipitation response to GHG forcings is smaller per unit change in forcing, than it is for short wave radiative forcings like volcanic aerosol¹⁴. Overall anthropogenic forcings result in a lower rate of increase in precipitation globally than suggested by the Clausius-Clapeyron relation^{14, 13,15,20,21,22}.

A pioneering study¹⁴ quantified the expected range of change in total global precipitation in response to CO₂ driven warming, but found that even at large scales there was considerable variation in the expected spatial pattern of change. A key

advance in the physical explanation of the response pattern of precipitation changes due to increasing GHGs was made by a later study¹⁵. They identified robust features of anthropogenic changes such as enhancement of the patterns of precipitation minus evaporation (P-E), poleward movement of the Hadley circulation and subsequent shifting of the arid subtropical subsidence regions and storm tracks, leading to the ‘wet gets wetter’ and ‘dry gets drier’ paradigm. It has recently been found that although this paradigm has some validity over wet higher latitudes and dry subtropical land regions, it does not hold true everywhere. For example, humid to transitional regimes are shifting to drier conditions²³. Other changes in large-scale rainfall patterns have been explained through a ‘warmer-get-wetter’ mechanism, by which warm SST patterns over the tropics cause increases in precipitation²⁴.

Expectation of regional changes

Change in regional rainfall is a consequence both of thermodynamics and anthropogenic influence on dynamics²⁵. Human-induced depletion in stratospheric ozone, for example, is found to cause a poleward shift of the southern extratropical jets, which affect regional precipitation patterns in the Southern Hemisphere^{26, 27}. The storm track in the Northern Hemisphere, and hence rainfall in Europe, are also affected by changes in stratospheric circulation²⁸.

More generally, the regional precipitation response to naturally occurring modes of variability, such as ENSO and the NAO, is influenced by the basic state of the atmosphere and ocean^{14, 29,30}. It is to be expected therefore that anthropogenic perturbations to the basic state would lead to changes in regional teleconnection patterns.

The regional character of anthropogenic precipitation change, therefore, results from complex interactions between natural variability and anthropogenic forcing. This is especially the case at regional scales. Indeed, variability related to teleconnections is not likely to affect total precipitation over very large domains, because wetter conditions in one place tend to be balanced by dryer conditions elsewhere³¹. In short, in order to disentangle the complex causes of regional precipitation change, we need to consider the following three aspects of the response: 1) external forcing may project onto internal variability, changing the amplitude or frequency of modes of climate variability, or altering the teleconnections that govern precipitation response, 2) the fingerprint of external forcing may reflect both thermodynamic and dynamic changes through forced changes to atmospheric energetics, moisture content, and large-scale circulation, and 3) the precipitation responses to different external drivers such as greenhouse gases, aerosols, ozone, natural events will differ.

Modelling and observational uncertainties

Recent studies that have sought to detect and attribute anthropogenic signals in large-scale zonal precipitation have compared observations to CMIP5 (Coupled Model Intercomparison Project 5) model simulations with and without anthropogenic forcings^{2, 3}. Anthropogenic increases in precipitation on global land and ocean are clear in model simulations (Figure 1a-c). However attribution approaches require that like is compared with like by comparing observations of the historical period to models that have been masked with the observational coverage. This means that the clear signals seen in models are obscured by sparse observational coverage². These findings indicate that global as well as zonal trends are distorted by the aliasing of sparse observational coverage onto the multi-model means.

The robustness of the detection of global and large-scale trends (Figures 10.10 & 10.A.2 of ref. 6) needs to be tested by comparing model data with different datasets of long-term observations. Ref. 2, for example, detected seasonal changes in zonal-mean precipitation attributable to human activities in four observational datasets – albeit only for March-April-May and December-January-February. However, the magnitudes of the temporal fingerprint of mid-to-high latitude positive trends and low latitude negative trends vary between observational datasets (Figure 2). In fact, anthropogenic changes are detected for all seasons in only one of the observational datasets³. The sensitivity of findings to observational dataset illustrates the barriers imposed by observational uncertainty.

The above discussion has focussed on uncertainties in observations of precipitation. It should not be forgotten, however, that effective model-observation comparison relies on accurate observations, not only of the variable in question, but also of forcing factors, including natural and anthropogenic aerosol. It has been found, for example, that natural desert dust aerosols from North Africa and West Asia are positively correlated to Indian summer monsoon rainfall on short time scales, with the dust-induced heating favouring increased moisture convergence over the Arabian peninsula and hence the westerly flow and precipitation over the Indian subcontinent³². Such model based findings point to the increasing need for an improved understanding of the climatic response to aerosols, which will require more systematic modelling experiments exploring the sensitivity of the precipitation response to aerosol forcing uncertainty as well as improvements in the representation of aerosol forcing in models.

Many of the impacts of a changing water cycle are felt at regional and local scales rather than at continental or global scales. Observational uncertainty at any given grid

point (of resolution of a few hundreds of kms) may be greatest at these scales (http://sciforum.net/conference/66/paper/2901). Paradoxically, however, observational uncertainty may be less of a barrier to attribution at the regional than at the global level. At the largest spatial scales, many of the detection and attribution issues related to observational uncertainty stem from sparse spatial sampling² in observations which means that the trends from models and observations can be badly distorted, losing much of the underlying signals. At local scales, in contrast, inconsistency in spatial sampling is less likely to contribute significantly to observational uncertainty. Instead, observational uncertainty reflects the sparsity of ground observations and consequent measurement/calibration errors. Such uncertainty may not, in itself, preclude robust detection and attribution of anthropogenic change in some regions, providing there exist temporally consistent ground or satellite based rainfall estimates. Indeed, at these scales, detection and attribution may be hampered more by the challenge of comparing models and observations, than by observational uncertainty itself. This is, in part because there are large discrepancies between the locations of simulated and observed features in the climatologies of precipitation which might be expected to cause differences in the anthropogenic response³³. These discrepancies are compounded by the lack of robustness in model-simulated internal variability³⁴ causing uncertainty in the fingerprint^{3, 35}, or under sampling of the observed variability³⁶ – which as described in earlier sections are a particularly serious issue at the regional scale.

A clearer view

The success of any approach to detection and attribution is contingent on the model's ability to represent the relevant processes over a particular region and season. Structural uncertainties in climate models (due to the differences in models' structure

leading to individual model errors), although reduced since the Fourth Assessment Report^{37, 38} (AR4), remain as a barrier to quantifying robust change in precipitation on regional scales³⁹.

The need for improved process-representation has motivated recent work on improved model dynamics and resolution⁴⁰, and the incorporation of individual processes and complex models of individual parts of the climate system⁴¹. High horizontal and vertical resolution and improved parameterisations in climate models have been shown to improve representation in models of processes, such as the vorticity of tropical cyclones, storm dynamics, atmospheric fronts, convection and blocking, clouds and their interactions with aerosols, gravity waves, ocean-biogeochemistry, land and sea-ice, boundary layer and land-surface processes, and strength of the local hydrological cycle^{40, 41, 42, 43, 44, 45}. The development of both high-resolution climate models and Earth System Models (ESMs) are thus instrumental in tackling regional climate problems. Ref. 40, for example, performed climate change experiments using a 1.5 km resolution regional climate model and projected future increase in heavy downpours over the UK. They illustrated that explicit convection and local storm dynamics are important in simulating the fine temporal and spatial scales of UK summer rainfall.

Compared to CMIP3 models, many CMIP5 models represent first and second indirect effect of aerosols and improved aerosol-cloud representations. On large spatial scales, these significant improvements in climate model representation of aerosols have now enabled improved simulation of inter-decadal variability in temperature and precipitation^{35, 46}. A weakening of the Northern Hemisphere land precipitation between the 1950s and 1980s and a subsequent recovery has been detected and attributed to increasing anthropogenic aerosols during 1950 to 1980s followed by a re-

emergence of the greenhouse gas signal relative to the anthropogenic aerosol signal in later years³⁵. Models with representation of the indirect effect of sulphate aerosols, together with the direct effect of sulphate aerosols perform better in representing the rate of decrease of precipitation in the 1950s and the recovery in the 1980s than the models that exclude the indirect effect⁴⁶ although models still have shortcomings in representing the timing of the recovery. There is thus a scientific opportunity to use these newly available simulations to decipher the joint influence of anthropogenic aerosols and greenhouse gas emissions on regional precipitation, and hence to detect anthropogenic trends.

New methodologies

The base climate is expected to vary from one model to another. Averaging simplistically over output from many models may therefore obscure signals of anthropogenic change. For instance, variation between models of the location and seasonal timing of precipitation may hamper robust assessment of changes in the mean^{33, 47,48}. Novel methods of accounting for the mismatches between model climatologies offer a means of tackling the problem of consistent model changes being distorted by differences in climatological features (eg. convergence zones) both between models, and between models and observations^{33,49}. In order to correct feature location errors in GCMs, ref. 33 applied a warping method, which has been used in brain imagery registration, to monthly precipitation fields. The warping technique was found to improve the detectability of human influence⁴⁹. Other model-observation comparison methods such as the model-by-model approach⁴⁸ and space-scale smoothing⁴⁷, which consider individual model runs as opposed to the multi-model ensemble mean, have also been shown to reduce feature-location biases and hence to identify robust changes in the location and magnitude of zonal extremes.

Natural variability, as well as systematic bias in models, can obscure part of the signal of anthropogenic change in precipitation. For example, the anthropogenic effect on the precipitation response to natural modes of variability is superposed on natural variation in the amplitude and frequency of these modes^{50, 51,52,53}. Aliasing natural internal variability and changes due to anthropogenic forcing in this manner would be expected to cause variations in the anthropogenic effect on regional precipitation. So if, say, greenhouse gas forcing modifies the precipitation response to ENSO in a given region, the anthropogenic expression of precipitation change is more pronounced during periods when ENSO is active. These periods cannot be expected to coincide in free-running coupled climate models. Averaging precipitation over large model ensembles will therefore not reveal this component of the signal of anthropogenic influence. Rather detection and attribution techniques need to take explicit account of the drivers of precipitation variability (e.g. ENSO, NAO) and to their effects on precipitation (e.g. ENSO teleconnections) rather than just treating such variability as noise in the analysis. This type of process-based approach complements the application of detection and attribution techniques directly to regional precipitation^{8,9} and can yield a clearer understanding of the role of natural and anthropogenic factors⁷¹.

On regional scales, therefore, in addition to analysing precipitation directly, it is productive to investigate the processes underlying precipitation change (process-based fingerprints). Examples of such fingerprints are the increased risk of heavy rainfall during mid-latitude atmospheric river events in the UK^{54, 55} and New Zealand⁵⁶; the poleward migration of the storm track⁴⁷ (Figure 3) and the large scale dynamical implications of an expected intensification of the hydrological cycle^{15, 20, 57,58} that, at least over non-water limited regions²³ of the earth including the oceans, many wet

regions tend to get wetter and dry regions drier. As pointed out earlier it should be noted that the over simplicity of this expectation from theory and models is currently under discussion²³. However, a temporal response pattern with wet tropical regions getting wetter and dry regions getting drier was apparent in simulations of the recent past and future projections from CMIP5 models and was consistent with satellite rainfall observations for the tropical region²⁰. ENSO variability can cause increase or decrease of regional rainfall over the land depending on the sign of the phase⁵⁸ suggesting if the ENSO characteristics change such precipitation response which is governed by remote SST patterns may change too. On fine scales, shifting of the wet and dry regions may make it difficult to identify this expected pattern of change^{23, 59,60}. However, using two fingerprints of wet and dry processes, ref. 57 detected an expected intensification of the water cycle partly attributable to human-induced greenhouse gas forcing.

Anthropogenic change in precipitation is driven not only by greenhouse gas emission, but also by aerosol forcing which modulates regional precipitation. Sulphate aerosol and desert dust forcings influence changes in the wet and dry conditions of Sahelian water cycle caused primarily by changes in West African Monsoon rains through changes in SST feedbacks and subsequent shifts in tropical convergence zones^{61, 62}. Simulated Sahel rainfall is found to weaken due to rapid changes in anthropogenic sulphur dioxide emissions from Asia and Europe through a fast (less than 3 weeks) aerosol-radiation and aerosol-cloud response and a slow (more than 3 weeks) response (i.e. decrease in West African Monsoon by adjustment of Walker circulation) caused by atmosphere and land-surface feedbacks⁶³. While there was a decrease of Sahel rainfall during the 1970s and 1980s since then there has been some recovery of Sahel rainfall which could have been influenced by increasing levels of

greenhouse gases in the atmosphere as well as changes in anthropogenic aerosol precursor emissions⁶⁴.

Event attribution

The previous discussion has highlighted the importance of identifying and isolating processes underlying anthropogenic change in precipitation. This can be accomplished, as described in the studies cited above, by explicitly isolating candidate processes and investigating how they are affected by anthropogenic climate change. A further refinement is to investigate the anthropogenic contribution to the processes underpinning individual extreme events – a technique known as event attribution.

Event attribution studies seek to determine how anthropogenic forcings have altered the magnitude or probability of a particular type of extreme weather or climate-related event as experienced in the observed record^{65, 66, 67}. In recent years efforts have been made to carry out such studies shortly after the events in question, for example in the publication of an annual series of reports which explain extreme events of the previous year from a climate perspective⁶⁸. However while there is increasing evidence that robust attribution statements can be made about an anthropogenic contribution to the likelihood of many extreme warm events, the role of human influences on extreme precipitation events is decidedly mixed⁶⁹ consistent with previous findings about the difficulties of robustly attributing precipitation events. Nevertheless such diagnostic approaches to attribution have made some headway in breaking down the problem into thermodynamic and dynamical components⁷⁰ and in devising modelling strategies to quantify the different contributions from anthropogenic and natural forcings and aspects of internal variability⁶⁴. It is therefore becoming possible to attribute changes in probability of some types of regional

extreme precipitation event through developing an understanding of the thermodynamic and dynamic contributors^{71, 72}. Ref. 73 argues that in attributing extreme climate events it is more useful to regard the extreme circulation regime or weather event as being largely unaffected by climate change and to concentrate solely on the thermodynamic component of an anthropogenic impact on the event in question. However it is important to consider dynamic factors as well as thermodynamic factors and to consider the extent to which dynamical aspects may have changed since it is both that contribute to the risk of a particular event^{74, 71, 72, 75}. Also attention should be given as to whether there are non-linear interactions between the two, as discussed above.

The way ahead

Based on our discussion of scientific opportunities and challenges, we emphasise that quantification of the effects of human influence on precipitation across the globe crucially depends on developing and applying process understanding. Given current observational uncertainties⁴ and limitations in models³⁸ simply waiting for improvements in observations and models to deliver clearer detection and attribution results seems an insufficient response to the challenge of better understanding how climate change is affecting precipitation around the globe. For example some of the important recommendations proposed by ref. 4 such as the observational data rescue, improvements in the observational coverage and models could take years to implement. Clearly observations and models are continuously improving and detection and attribution analyses should take advantage of such advances. But adaptation decisions could be even better informed if it were possible to incorporate process understanding more in detection and attribution studies. Those adaptation decisions that are based on robust climate projections are much stronger where the

projections are based on firm foundation of physical understanding and underpinned by robust attribution studies. Hence attribution studies are central to informed adaptation planning and decision making. Even where large uncertainties remain, additional useful information could be obtained and applied in a risk-based framework⁶⁰ based on an understanding of the likely mechanisms at work.

In particular, we need to better understand the expected effect of anthropogenic climate change on modes of variability and their teleconnections with regional precipitation²⁹. Disentangling these effects will allow an improved understanding of the extent to which regional changes are anthropogenically caused versus being caused by natural variations, either internally generated within the climate system or externally forced, such as by solar variability or explosive volcanic eruptions. It is not always reasonable to consider internal variability simply as ‘noise’ to be filtered out.

Recent process-based detection and attribution approaches⁴⁷, which consider the signal or the forced response being thermodynamic and/or dynamic in origin, have shown some success. There is indication that the anthropogenic signal could also be expressed in part through changes in amplitude, frequency and modes of natural internal variability. An alternative approach would be to look directly at the anthropogenic signal as a net effect of rainfall changes due to a) thermodynamic contribution, b) dynamic contribution (which includes changes in circulation, modes of variability and changes in teleconnections due to changes in modes of variability). Analyses quantifying changes in natural internal variability⁷⁶ would be a valuable addition to quantifying forced changes over regions where internal variability on interannual timescales is changing. However, it is very difficult to robustly detect changes in observed variability for a highly noisy climate variable as precipitation.

New metrics that best express robust changes in the water cycle would aid in identifying anthropogenic changes. For example this could involve calculating areas of land with precipitation changes at particular thresholds¹² or could involve combining terrestrial observations of precipitation with oceanographic observations of salinity⁶.

In summary, we have shown that, even in the face of imperfect models and observations, progress can be made in detecting and attributing changes in regional precipitation. Improved process understanding, innovations in detection and attribution methodologies, and novel methods of confronting models with observations can now be brought to bear on this highly challenging problem. Development of high quality observational datasets and high-resolution models will be undoubtedly helpful and are likely to have substantial pay off over the longer term. But in the meantime, innovative methods for analysing the observations and models we have available now could yield important additional information to inform societies and policy makers about the nature of changing precipitation at fine spatial-scales.

Box 1. What is detection and attribution?

Detection of a change is the process of demonstrating that climate has changed in some defined statistical sense, without providing a reason for that change⁷⁷.

Attribution of causes of the change is defined as the process of evaluating the relative contributions of multiple causal factors to a change or event with an assignment of statistical confidence⁶. *Fingerprints* are metrics or space-time patterns of the response of climate variables to anthropogenic forcings, such as greenhouse gas emissions, atmospheric pollutants, or natural forcings such as solar radiation changes and

aerosols from explosive volcanic eruptions. Most of the recent detection and attribution studies use climate models⁷⁸ to estimate the expected fingerprints of change and the uncertainty of their estimate in observations of the real world. For an overview of techniques, see Appendix 9.2 of AR4⁶² and Section 10.2.1 of AR5⁶.

References:

1. IPCC, Summary for Policymakers. In: *Climate Change 2014: Impacts, Adaptation, and Vulnerability*. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (eds Field, C. B. *et al.*). Cambridge University Press. 1-32 (2014).
2. Balan Sarojini, B., Stott, P.A., Black, E. & Polson, D. Fingerprints of changes in annual and seasonal precipitation from CMIP5 models over land and ocean. *Geophys. Res. Lett.* **39**, L21706, DOI: 10.1029/2012GL053373 (2012).
3. Polson, D., Hegerl, G.C., Zhang, X. & Osborn, T. Changes in seasonal land precipitation during the latter twentieth-century. *J. Clim.* **20**, 6679-6697 (2013a).
4. Hegerl, G.C. *et al.* Challenges in quantifying changes in the global watercycle. *Bull. Am. Meteorol. Soc.* DOI:10.1175/bams-d-13-00212.1 (2015).
5. IPCC, *Climate Change 2013: The Physical Science Basis*. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (eds Stocker, T.F., *et al.*). Cambridge University Press (2013).
6. Bindoff, N.L. *et al.* Detection and Attribution of Climate Change: from Global to Regional. In: *Climate Change 2013: The Physical Science Basis*. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel

- 454 on Climate Change [eds Stocker, T.F. *et al.*]. Cambridge University Press, 867-
455 952 (2013).
- 456 7. Noake, K., Polson, D., Hegerl, G. & Zhang, X. Changes in seasonal land
457 precipitation during the latter twentieth-century. *Geophys. Res. Lett.* **39**, L03706
458 (2012).
- 459 8. Min, S., Zhang, X. & Zwiers, F.W. Human-induced Arctic moistening. *Science*
460 **320**, 518-520 (2008).
- 461 9. Wan, H, *et al.* Attributing northern high-latitude precipitation change over the
462 period 1966-2005 to human influence. *Clim. Dyn.*, DOI: 10.1007/s00382-014-
463 2423-y (2014).
- 464 10. Delworth, T.L. & Zeng, F. Regional rainfall decline in Australia attributed to
465 anthropogenic greenhouse gases and ozone levels. *Nature Geosci.* **7**, 583–587,
466 DOI: 10.1038/ngeo2201 (2014).
- 467 11. Zhang, X, *et al.* Detection of human influence on twentieth-century precipitation
468 trends. *Nature* **448**, 461–465 (2007).
- 469 12. Chadwick, R., Good P., Martin G. & Rowell D.P. Large rainfall changes
470 consistently projected over substantial areas of tropical land. *Nature Clim.*
471 *Change*. **5**, 1-5 (2015).
- 472 13. Trenberth, K.E. Changes in precipitation with climate change. *Clim. Res.* **47**, 123–
473 138 (2011).
- 474 14. Allen, M.R., & Ingram, W.J. Constraints on future changes in climate and the
475 hydrologic cycle. *Nature* **419**, 224–232, DOI: 10.1038/nature01092 (2002).
- 476 15. Held, I.M., & Soden, B.J. Robust responses of the hydrological cycle to global
477 warming. *J. Clim.*, **19**, 5686–5699 (2006).

- 478 16. Willett, K.M., Jones, P.D., Gillett, N.P. & Thorne, P.W. Attribution of observed
479 surface humidity changes to human influence. *Nature* **449**, 710-713,
480 DOI:10.1038/nature06207 (2007).
- 481 17. Santer, B.D. *et al.* Identification of human-induced changes in atmospheric
482 moisture content. *Proc. Natl. Acad. Sci.* **104**, 15248-15253 (2007).
- 483 18. Santer, B.D. *et al.* Incorporating model quality information in climate change
484 detection and attribution studies. *Proc. Natl. Acad. Sci.* **106**, 14778-14783 (2009).
- 485 19. Blunden, J., and Arndt, D.S. Eds., State of the Climate in 2013. *Bull. Amer.*
486 *Meteor. Soc.* **95**, 1-238 (2014).
- 487 20. Allan, R.P. *et al.* Physically consistent responses of the global atmospheric
488 hydrological cycle in models and observations. *Surv. Geophys.* DOI:
489 10.1007/s10712-012-9213-z (2013).
- 490 21. Pendergrass, A.G. & Hartmann, D.L. The atmospheric energy constraint on
491 global-mean precipitation change. *J. Clim.*, **27**, 757–768 (2014).
- 492 22. Thorpe, L. & Andrews, T. The physical drivers of historical and 21st century
493 global precipitation changes. *Environ. Res. Lett.* **09**, 064024 (2014).
- 494 23. Greve, P. *et al.* *Nature Geosci.* **7**, 716–721 (2014).
- 495 24. Xie, S.-P. *et al.* Global warming pattern formation: Sea surface temperature and
496 rainfall. *J. Clim.*, **23**, 966-986 (2010).
- 497 25. Seager, R.J. Thermodynamic and dynamic mechanisms for large-scale changes in
498 the hydrological cycle in response to global warming. *J. Clim.*, **23**, 4651-4668
499 (2010).
- 500 26. Kang, S.M., Polvani, L.M., Fyfe, J.C. & Sigmond, M. Impact of polar ozone
501 depletion on subtropical precipitation. *Science* **332**, 951–954 (2011).

- 502 27. Min, S.-K., & Son, S.-W., Multimodel attribution of the Southern Hemisphere
503 Hadley cell widening: Major role of ozone depletion, *J. Geophys. Res.*
504 *Atmos.*, **118**, 3007–3015, DOI:10.1002/jgrd.50232 (2013).
- 505 28. Scaife, A. *et al.* Climate change projections and stratosphere–troposphere
506 interaction, *Clim. Dyn.* **38**, 2089–2097 (2012).
- 507 29. Seager, R.J., Naik, N & Vogel, L. Does Global Warming Cause Intensified
508 Interannual Hydroclimate Variability? *J. Clim.*, **25**, 3355–3372 (2012).
- 509 30. Vecchi, G.A., & Wittenberg, A.T., El Niño and our future climate: where do we
510 stand? Wiley Interdisciplinary Reviews: *Climate Change* **1.2**, 260-270 (2010).
- 511 31. Nicholson, S.E., & Kim, J. The relationship of the El Nino-Southern oscillation to
512 African rainfall. *International Jn. of Clim.* **17.2**, 117-135 (1997).
- 513 32. Vиноj, V. *et al.* Short-term modulation of Indian summer monsoon rainfall by
514 West Asian dust. *Nature Geosci.* **7**, DOI: 10.1038/ngeo2107 (2014).
- 515 33. Levy, A.A.L. *et al.* Can correcting feature location in simulated mean climate
516 improve agreement on projected changes? *Geophys. Res. Lett.* **40**, DOI:
517 10.1029/2012GL053964 (2013).
- 518 34. Collins, M. *et al.* Observational challenges in evaluating climate models. *Nature*
519 *Clim. Change* **3**, 940-941 (2013).
- 520 35. Wu, P., Christidis, N. & Stott, P.A. Anthropogenic impact on Earth's hydrological
521 cycle. *Nature Clim. Change* DOI: 10.1038/NCLIMATE1932 (2013).
- 522 36. Wan, H. *et al.* Effect of data coverage on the estimation of mean and variability of
523 precipitation at global and regional scales, *J. Geophys. Res. Atmos.*, **118**, 534–546,
524 DOI:10.1002/jgrd.50118 (2013).

- 525 37. Flato, G. *et al.* Evaluation of Climate Models. In: *Climate Change 2013: The*
526 *Physical Science Basis*. Contribution of Working Group I to the Fifth Assessment
527 Report of the Intergovernmental Panel on Climate Change [eds Stocker, T.F. *et*
528 *al.*]. Cambridge University Press. (2013).
- 529 38. Stevens, B. & Bony, S. What are climate models missing? *Science* **340**, 1053–
530 1054 (2013).
- 531 39. Knutti, R. & Sedlacek, J. Robustness and uncertainties in the new CMIP5 climate
532 model projections. *Nature Clim. Change* **3**, 369–373 (2013).
- 533 40. Kendon, E.J. *et al.* Heavier summer downpours with climate change revealed by
534 weather forecast resolution model. *Nature Clim. Change* **4**, 570–576, DOI:
535 10.1038/nclimate2258 (2014).
- 536 41. Cox, P.M. *et al.* Sensitivity of tropical carbon to climate change constrained by
537 carbon dioxide variability. *Nature* **494**, 341–344, DOI: 10.1038/nature11882
538 (2013).
- 539 42. Roberts, M.J. *et al.* Tropical cyclones in the UPSCALE ensemble of high
540 resolution global climate models. *J. Clim.*, **28**, 574–596. ISSN 1520-0442
541 DOI: 10.1175/JCLI-D-14-00131.1 (2015).
- 542 43. Demory, M.-E. *et al.* The role of horizontal resolution in simulating drivers of the
543 global hydrological cycle. *Clim. Dyn.*, **42**, 2201–2225. ISSN 0930-7575
544 DOI: 10.1007/s00382-013-1924-4 (2013).
- 545 44. Jung, T., *et al.* High-Resolution Global Climate Simulations with the ECMWF
546 Model in Project Athena: Experimental Design, Model Climate, and Seasonal
547 Forecast Skill. *J. Clim.*, **25**, 3155–3172. DOI:10.1175/JCLI-D-11-00265.1 (2012).

- 548 45. Strachan, J., Vidale, P.L., Hodges, K., Roberts, M. and Demory, M.-
549 E. Investigating global tropical cyclone activity with a hierarchy of AGCMs: the
550 role of model resolution. *J. Clim.*, **26**, 133-152. ISSN 1520-0442
551 DOI: 10.1175/JCLI-D-12-00012.1 (2013).
- 552 46. Wilcox, L.J., Highwood, E.J. & Dunstone, N.J. The influence of anthropogenic
553 aerosol on multi-decadal variations of historical global climate. *Environ. Res. Lett.*
554 **8** (2). 024033. ISSN 1748-9326 DOI: 10.1088/1748-9326/8/2/024033 (2013).
- 555 47. Marvel, K. & Bonfils, C. Identifying external influences on global precipitation.
556 *Proc. Natl. Acad. Sci.* www.pnas.org/cgi/DOI/10.1073/pnas.1314382110 (2013).
- 557 48. Scheff, J & Frierson, D. Twenty-First-Century Multimodel Subtropical
558 Precipitation Declines Are Mostly Midlatitude Shifts. *J. Clim.*, **25**, 4330–4347
559 (2012).
- 560 49. Levy, A.A.L. *et al.* Correcting feature location in GCMs aids the detectability of
561 external influence on precipitation. *J. Geophys. Res.*, (2014).
- 562 50. Corti, S., Molteni, F. & Palmer, T.N., Signature of recent climate change in
563 frequencies of natural atmospheric circulation regimes, *Nature*, 398, 799–802,
564 (1999).
- 565 51. Mann, M.E., Bradley, R.S. & Hughes, M.K., Long-term variability in the ENSO
566 and associated teleconnections, in *ENSO: Multiscale Variability and Global and*
567 *Regional Impacts*, edited by H.F. Diaz and V. Markgraf, pp. 357–412, Cambridge
568 Univ. Press, New York. (2000).
- 569 52. Black, E. The influence of the North Atlantic Oscillation and European circulation
570 regimes on the daily to interannual variability of winter precipitation in
571 Israel. *International Jn. of Clim.*, DOI: 10.1002/joc.2383 (2011).

53. Christensen, J.H. *et al.* Climate Phenomena and their Relevance for Future Regional Climate Change. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (Stocker, T.F. *et al.* (eds.)). Cambridge University Press. (2013).
54. Lavers, D.A. *et al.* The detection of atmospheric rivers in atmospheric reanalyses and their links to British winter floods and the large-scale climatic circulation. *J. Geophys. Res.* **117**, D20106, DOI: 10.1029/2012JD018027 (2012).
55. Lavers, D.A. *et al.* Future changes in atmospheric rivers and their implications for winter flooding in Britain. *Environ. Res. Lett.* **8**, 034010, DOI:10.1088/1748-9326/8/3/034010 (2013).
56. Dean, S.M., Rosier, S., Carey-Smith, T. & Stott, P.A. The role of climate change in the two-day extreme rainfall in Golden Bay, New Zealand, December. [In “Explaining Extreme Events of 2012 from a Climate Perspective”]. *Bull. Amer. Meteorol. Soc.* **94** (9), S61–S63, DOI:10.1175/BAMS-D-13-00212.1 (2013).
57. Polson, D., Hegerl, G.C., Allan, R.P. & Balan Sarojini, B. Have greenhouse gases intensified the contrast between wet and dry regions? *Geophys. Res. Lett.* **40**, DOI:10.1002/grl.50923 (2013b).
58. Liu, C. & Allan, R.P. Observed and simulated precipitation responses in wet and dry regions 1850–2100. *Environ. Res. Lett.* **8**, 034002, DOI:10.1088/1748-9326/8/3/034002 (2013).
59. Allan, R.P. Climate Change: Dichotomy of drought and deluge. *Nature Geosci.* DOI: 10.1038/ngeo2243 (2014).

- 595 60. Shepherd, T.G. Atmospheric circulation as a source of uncertainty in climate
596 change projections. *Nature Geoscience*. DOI: 10.1038/ngeo2253, (2014).
- 597 61. Rotstayn, L.D., & Lohmann, U. Tropical rainfall trends and the indirect aerosol
598 effect. *J. Clim.*, **15**, 2103–2116 (2002).
- 599 62. Hegerl, G.C. *et al.* Understanding and Attributing Climate Change. In: *Climate*
600 *Change 2007: The Physical Science Basis*. Contribution of Working Group I to
601 the Fourth Assessment Report of the Intergovernmental Panel on Climate Change
602 (Solomon, S. *et al.* (eds.)). Cambridge University Press. (2007).
- 603 63. Dong, B., Sutton, R., Highwood, E.J. & Wilcox, L.J. The impacts of European
604 and Asian anthropogenic sulphur dioxide emissions on Sahel rainfall. *J. Clim.*, **27**,
605 7000-7017 (2014).
- 606 64. Dong, B.-W. & Sutton, R., Dominant role of greenhouse gas forcing in the
607 recovery of Sahel rainfall. *Nature Clim. Change*. DOI: 10.1038/nclimate2664
608 (2015).
- 609 65. Allen, M.R. Liability for climate change. *Nature* **421**, 891–892 (2003).
- 610 66. Stott, P.A., Stone, D.A., & Allen, M.R. Human contribution to the European heat
611 wave of 2003. *Nature*, **432**, 610–614 (2004).
- 612 67. Pall, P., *et al.*, Anthropogenic greenhouse gas contribution to UK autumn flood
613 risk. *Nature*, **470**, 382–385 (2011).
- 614 68. Herring, S.C., Hoerling, M.P., Kossin, J.P., Peterson, T.C. & Stott, P.A., Eds.,
615 Explaining Extreme Events of 2014 from a Climate Perspective. *Bull. Amer.*
616 *Meteor. Soc.*, **96** (12), S1–S172 (2015).
- 617 69. Herring, S., *et al.* Summary and Broader context [in "Explaining Extremes of
618 2013 from a Climate Perspective"]. *Bull. Amer. Meteor. Soc.*, **82** (9) (2014b).

- 619 70. Hoerling, M., *et al.* Northeast Colorado extreme rains interpreted in a climate
620 change context [in "Explaining Extremes of 2013 from a Climate Perspective"].
621 *Bull. Amer. Meteor. Soc.*, **95** (9), S15–S18 (2014).
- 622 71. Christidis, N.C. & Stott, P.A. Extreme rainfall in the United Kingdom during
623 winter 2013/14: the role of atmospheric circulation and climate change [in
624 "Explaining Extremes of 2014 from a Climate Perspective"]. *Bull. Amer. Meteor.*
625 *Soc.*, **96** (12), S46–S50 (2015).
- 626 72. Schaller, N., *et al.*, Human influence on climate in the 2014 Southern England
627 winter floods and their impacts. *Nature Clim. Change*. (2016).
- 628 73. Trenberth *et al.* Attribution of climate extreme events. *Nature Clim. Change*.
629 DOI: 10.1038/NCLIMATE2657 (2015).
- 630 74. Huntingford C., *et al.* Potential influences in the United Kingdom's floods of
631 winter 2013–2014. *Nature Clim. Change*, **4**, 769–777 (2014).
- 632 75. Stott, P.A., *et al.*, Attribution of extreme weather and climate-related events.
633 *WIREs Clim. Change*, **7**, 23–41. DOI: 10.1002/wcc.380 (2016).
- 634 76. Power *et al.* Robust twenty-first-century projections of El Niño and related
635 precipitation variability. *Nature*. **502**, 541–545 (2013).
- 636 77. Hegerl, G.C. *et al.* Good practice guidance paper on detection and attribution
637 related to anthropogenic climate change. In: *Meeting Report of the*
638 *Intergovernmental Panel on Climate Change Expert Meeting on Detection and*
639 *Attribution of Anthropogenic Climate Change* (eds Stocker, T.F. *et al.*). IPCC
640 Working Group I Technical Support Unit. (2010).
- 641 78. Hegerl, G.C. & Zwiers, F.W. Use of models in detection and attribution of climate
642 change. *WIREs Clim. Change*, **2**, 570–591 (2011).

79. Harris, I. et al. Updated high-resolution grids of monthly climatic observations - the CRU TS 3.1 Dataset. *International Jn. of Clim.* **34**, 623-642 (2014).

80. Becker, A. et al. A description of the global land-surface precipitation data products of the Global Precipitation Climatology Centre with sample applications including centennial (trend) analysis from 1901-present. *Earth Syst. Sci. Data Discuss.*, **5**, 971-998. DOI: 10.5194/essd-5-71-2013 (2013).

81. Beck, C., Grieser, J., & Rudolf, B. A new monthly precipitation climatology for the global land areas for the period 1951 to 2000, *Climate Status Report*, 181-190 (2004).

Acknowledgements

This work is supported by Horyuji PAGODA project of the Changing Water Cycle programme of UK Natural Environment Research Council (NERC) (Grant NE/I006672/1) and by the Joint DECC/Defra Met Office Hadley Centre Climate Programme (GA01101). B.B.S. acknowledges joint support from the UK NERC (NE/I006672/1) and the Met Office Hadley Centre, and a discussion with Pier Luigi Vidale and Anne Verhoef on the atmospheric-land surface processes. E.B. was supported by the National Centre for Atmospheric Science – Climate division core research programme and the following research grants: HyCristal (NE/M020371/1), SatWIN-Scale (NE/M008797/1) and BRAVE (NE/M008983/1). The authors are thankful to three anonymous reviewers for their constructive and critical comments on the manuscript.

Author Contributions

B.B.S. developed the content and led the writing; P.A.S and E.B. designed the outline of the article, contributed to discussions, text, and commented on the drafts.

Competing Financial Interests statement

The authors declare no competing financial interests.

Corresponding Author

Correspondence to: Beena Balan Sarojini.

Figure Legends

Figure 1| Observational uncertainties due to sparse coverage obscure expected

fingerprints of change: Time-series of global mean precipitation anomalies (mm/day) w.r.t the baseline period of 1961-90, simulated by CMIP5 models forced with, both anthropogenic and natural forcings (ALL in orangish red lines) and natural forcings only (NAT in blue lines). a) Land and Ocean, b) Land, and c) Ocean, with all grid points. Multi-model means are shown in thick solid lines. Green stars show statistically significant changes at 5 % level. The clear signals seen in simulations (above) are obscured by sparse observational coverage when the global land precipitation is masked by observational coverage (Ref. 2).

Figure 2| Magnitudes of zonal mean land precipitation trends are dependent on

observational datasets: Comparison of observed trends (solid lines) using 4 observational datasets (Refs. 11, 79, 80, 81) for 1951-2005 (top). Range of CMIP5 simulations are in grey shading and multi-model ensemble mean (MM) in black dashed line. Blue (orange) shadings show latitudes where all observed datasets show positive (negative) trends. Comparison of simulated trends (bottom) using CMIP5 historical (ALL) simulations (individual simulations in grey dashed lines, multi-model mean in black dashed line), and the natural forcing only (NAT) simulations (MM in blue dashed lines) with the future (2006-2050) trend using RCP4.5 simulations (5-95 % range is in green shading, and MM in green dashed lines). Blue (orange) shading indicates latitudinal regions where more than two thirds of the historical simulations show positive (negative) trends (Ref. 4).

Figure 3| An example of simulated process-based fingerprint of anthropogenic precipitation change: Zonal mean boreal winter precipitation observations for 1990 (left). Local extrema are marked in dark blue (midlatitude storm tracks), red (subtropical dry zones), and green (equatorial tropical peak). Cyan, purple, and yellow circles indicate half-max points. Multivariate fingerprint $F_m(D,T)$ of forced precipitation change as thermodynamic (T) and dynamic (D) process indicators (right). Thermodynamic EOF loading is plotted on the vertical axis and the direction and magnitude of dynamic EOF loading are displayed as arrows showing the wet-gets-wetter and dry-gets-drier pattern in precipitation intensity and the poleward extension of precipitation over storm track and subtropical arid latitudes in both hemispheres (Ref. 47).

Figures




Figure 1| Observational uncertainties due to sparse coverage obscure expected fingerprints of change: Time-series of global mean precipitation anomalies

(mm/day) w.r.t the baseline period of 1961-90, simulated by CMIP5 models forced with, both anthropogenic and natural forcings (ALL in orangish red lines) and natural forcings only (NAT in blue lines). a) Land and Ocean, b) Land, and c) Ocean, with all grid points. Multi-model means are shown in thick solid lines. Green stars show statistically significant changes at 5 % level. The clear signals seen in simulations (above) are obscured by sparse observational coverage when the global land precipitation is masked by observational coverage (Ref. 2).




Figure 2| Magnitudes of zonal mean land precipitation trends are dependent on observational datasets: Comparison of observed trends (solid lines) using 4 observational datasets (Refs. 11, 79, 80, 81) for 1951-2005 (top). Range of CMIP5 simulations are in grey shading and multi-model ensemble mean (MM) in black dashed line. Blue (orange) shadings show latitudes where all observed datasets show positive (negative) trends. Comparison of simulated trends (bottom) using CMIP5 historical (ALL) simulations (individual simulations in grey dashed lines, multi-model mean in black dashed line), and the natural forcing only (NAT) simulations (MM in blue dashed lines) with the future (2006-2050) trend using RCP4.5 simulations (5-95 % range is in green shading, and MM in green dashed lines). Blue (orange) shading indicates latitudinal regions where more than two thirds of the historical simulations show positive (negative) trends (Ref. 4).




Figure 3| An example of simulated process-based fingerprint of anthropogenic precipitation change: Zonal mean boreal winter precipitation observations for 1990 (left). Local extrema are marked in dark blue (midlatitude storm tracks), red (subtropical dry zones), and green (equatorial tropical peak). Cyan, purple, and yellow

739 circles indicate half-max points. Multivariate fingerprint F_m (D, T) of forced
740 precipitation change as thermodynamic (T) and dynamic (D) process indicators
741 (right). Thermodynamic EOF loading is plotted on the vertical axis and the direction
742 and magnitude of dynamic EOF loading are displayed as arrows showing the wet-
743 gets-wetter and dry-gets-drier pattern in precipitation intensity and the poleward
744 extension of precipitation over storm track and subtropical arid latitudes in both
745 hemispheres (Ref. 47).

746
747